

Generation of Nonclassical States of Microwave Radiation via Single Photon Detection

Luke C.G. Govia,^{1,2} Emily J. Pritchett,¹ and Frank K. Wilhelm^{1,2}

¹*Theoretical Physics, Universität des Saarlandes, Saarbrücken, Germany*

²*Institute for Quantum Computing and Department of Physics and Astronomy, University of Waterloo, Ontario, Canada*

We describe the creation of nonclassical states of microwave radiation via ideal dichotomic single photon detection, i.e., a detector that only indicates presence or absence of photons. Ideally, such a detector has a back action in the form of the subtraction operator. Using the non-linearity of this back action, it is possible to create nonclassical states of microwave radiation, including squeezed and cat-like states, starting from a coherent state. We discuss the applicability of this protocol to current experimental designs of Josephson Photomultipliers (JPMs).

The generation of nonclassical states of light is an important test of the foundations of quantum mechanics and a necessary precursor to implementing quantum communication and computation protocols in many architectures [1–3]. While the methodology for creating nonclassical light at optical wavelengths has been studied extensively [4–6], the technology to create quantum states with larger and larger wavelengths has recently become available with advances in circuit-QED.

In this paper we present a novel way to generate nonclassical states of microwave radiation in a long wavelength transmission line using only detection by an ideal binary detector, the Josephson Photomultiplier (JPM). The protocol only involves radiating a microwave cavity with coherent light, and post selection based on single photon detection by JPMs without further manipulation. In addition, our protocol applies to any detection mechanism with a back action resembling that of the subtraction operator (equation (1)) and so can be generalized to other quantum systems, in particular, other superconducting circuits where strong photon-detector coupling is possible [7–9], but also at optical frequencies.

In the microwave regime, squeezed states [10–12] and cat-like states [13] of microwave radiation have been generated through the QND measurement of a microwave cavity with a Josephson parametric amplifier. We consider a projective detection scheme whereby the isolated cavity is left in a pure state, a protocol more similar to the Quantum Zeno detection scheme proposed for cavity-QED experiments [14]. We also note that the creation of arbitrary nonclassical states via the controlled interaction of a superconducting qubit with a microwave cavity has been achieved experimentally [8, 9]; here we show that a certain class of nonclassical states can be constructed more efficiently than thought previously.

The JPM, a current biased Josephson junction related to the phase qubit, has been shown experimentally [15] and theoretically [16–18] to be an effective single microwave photon counter. Previously, we have shown that for a JPM under optimal conditions the back action of photon detection is the photon subtraction operator [16],

$$\hat{B} \equiv \sum_{n=1}^{\infty} |n-1\rangle\langle n|, \quad (1)$$

a non-linear operator that can be related to the photon

lowering operator by $\hat{a} = \hat{B}\sqrt{\hat{n}}$, but cannot be expressed as a linear combination of photon creation and annihilation operators. Also, note that $\hat{B}$ is not invertible, and hence not unitary.

In this paper we show how to use the noncommutativity of the detection back action with coherent displacement pulses to achieve single mode quadrature squeezing of an input coherent state as well as other nonclassical states, including generalized Schrödinger cat states [4]. We also discuss the success probability of our protocol under realistic operating conditions.

The generation of squeezed states of microwave radiation using JPMs follows a simple protocol. The cavity is initially prepared in a coherent state, $|\alpha\rangle = e^{-\frac{|\alpha|^2}{2}} \sum_{n=0}^{\infty} \frac{\alpha^n}{\sqrt{n!}} |n\rangle$ where $\alpha \equiv |\alpha|e^{i\varphi_\alpha}$, and is coupled to one or more detectors, each acting with back action $\hat{B}$ on the cavity after a photon is detected. After N photons are counted, a displacement pulse is applied such that the state is centred around $-\alpha$ in phase space. After N further photon detections are observed, the resulting state is a squeezed state. The optimal choice of N will be discussed shortly.

Starting from the coherent state input, the probability for N photons to be detected is

$$P_N \equiv 1 - e^{-|\alpha|^2} \sum_{n=0}^{N-1} \frac{|\alpha|^{2n}}{n!} = 1 - \frac{\Gamma(N, |\alpha|^2)}{\Gamma(N)} \quad (2)$$

where $\Gamma(N, |\alpha|^2)$ is the upper incomplete gamma function of N and $|\alpha|^2$ [19]. $P_N \sim |\alpha|^{2N}/N!$ as $|\alpha| \rightarrow 0$; however, P_N increases monotonically with $|\alpha|$, so the success probability can be made arbitrarily close to unity with high power coherent pulses.

It is straightforward to calculate the normalized post measurement cavity state after N detections,

$$\rho' \equiv \frac{\hat{B}^N |\alpha\rangle\langle\alpha| \hat{B}^{\dagger N}}{P_N}, \quad (3)$$

and the average photon number $n' \equiv \langle a^\dagger a \rangle_{\rho'}$, is given by

$$n' = \frac{|\alpha|^2 \left(1 - \frac{\Gamma(N-1, |\alpha|^2)}{\Gamma(N-1)}\right) - N \left(1 - \frac{\Gamma(N, |\alpha|^2)}{\Gamma(N)}\right)}{P_N}, \quad (4)$$

or calculated straightforwardly numerically. After N detections, the next step is to displace the state by an amount $\alpha' = -\sqrt{n'}e^{i\varphi_\alpha} - \alpha$, so that the resulting state will be centred in phase space around $-\alpha$. For this displaced state, N photon detection events will occur with probability P'_N [27], and the renormalized cavity state will have the form

$$\rho'' \equiv \frac{\hat{B}^N D(\alpha') \rho' D(\alpha')^\dagger \hat{B}^{\dagger N}}{P'_N}. \quad (5)$$

We will now show that the state ρ'' is a squeezed state.

To quantify the amount of squeezing, we calculate the variance of the squeezed quadrature

$$\Delta p^2 = \text{Tr}[\rho'' \hat{p}^2] - \text{Tr}[\rho'' \hat{p}]^2. \quad (6)$$

The quadrature observable $\hat{p}(\phi_\alpha)$ is defined by $\hat{p}(\phi_\alpha) = \frac{1}{\sqrt{2}} (\hat{a}e^{-i\varphi_\alpha} - \hat{a}^\dagger e^{i\varphi_\alpha})$, where $\hat{a}$ is the annihilation operator for the cavity microwave mode. The phase shift $e^{-i\varphi_\alpha}$ accounts for the fact that this protocol squeezes along the phase space axes defined by the phase of the input coherent state. Anything less than $\Delta p^2 = \frac{1}{2}$ indicates a squeezed state. The amount of squeezing is expressed in dB, by calculating

$$S(\Delta p) \equiv 10\log_{10} \left(\frac{\Delta p^2}{\Delta p_{\text{norm}}^2} \right) = 10\log_{10} (2\Delta p^2). \quad (7)$$

In addition, we can calculate how far the state ρ'' deviates from a minimal uncertainty state by calculating $\Delta x \Delta p$ (where $\hat{x} = \frac{1}{\sqrt{2}} (\hat{a}e^{-i\varphi_\alpha} + \hat{a}^\dagger e^{i\varphi_\alpha})$ is the conjugate observable to $\hat{p}$). Figures 1(a) and 1(b) show $S(\Delta p)$ and $\Delta x \Delta p$ respectively as functions of α and the number of detection events on either side of the displacement, N .

As can be seen in figure 1(a), the maximum amount of squeezing possible on a given input state $|\alpha\rangle$ increases monotonically with $|\alpha|^2$, proportional to the power of the input pulse. Interestingly, for a given α there exists a finite N that achieves a global minimum in Δp^2 . One would be tempted to use this value of N in the protocol to create ρ'' , however, as can be seen in figure 1(b) there are other concerns.

As figure 1(b) shows, for a given α , $\Delta x \Delta p$ is not monotonic in N . Since we want ρ'' to be as close to a minimal uncertainty state as possible, while still maintaining a significantly squeezed quadrature, the optimal choice of N for the protocol would be at the $\Delta x \Delta p$ local minima shown in figures 1(a) and 1(b) by the white curve. While this does not minimize Δp^2 (and therefore maximize squeezing), it achieves a significantly squeezed state ρ'' that is as close to being a minimal uncertainty state as is possible, which is what we consider optimal.

Other nonclassical states of microwave radiation having $\varphi_k \equiv 2\pi/k$ rotational symmetry in phase space result from generalizing this procedure (see figure 2). The squeezed state protocol discussed previously involves detecting the cavity state while centred at two points on a line through the origin of phase space, i.e., the case




FIG. 1: (a) $S(\Delta p)$ and (b) $\Delta x \Delta p$ of the state ρ'' , as functions of α and the number of detection events, N . $\varphi_\alpha = 0$ for both figures. The white curve indicates the value of N that gives a local minimum in $\Delta x \Delta p$, while maintaining a squeezed Δp^2 .

$k = 2$. To generalize this protocol, we detect N photons at k positions equally spaced around a circle of radius $|\alpha|$ in phase space, and take the first position on the positive real axis for simplicity. There are k steps to this generalized protocol, and for $j = 0, 1, \dots, k-1$, the j th step is to 1) displace so that the cavity state is centred around $|\alpha|e^{ij\varphi_k}$, and 2) detect N photons. Finally, after detecting N photons at all k positions, we displace the state so that it's centred around the origin for comparison with the squeezed vacuum, and we refer to this final state as ρ_k . The detection stage at each position transforms ρ_{j1} to ρ_{j2} according to

$$\rho_{j2} \equiv \frac{\hat{B}^N \rho_{j1} \hat{B}^{\dagger N}}{P_N^j}, \quad (8)$$

where P_N^j is the probability of detecting N photons from the state ρ_{j1} . The entire protocol will complete with success probability

$$\text{Prob}(\text{success}) = \prod_{j=0}^{k-1} P_N^j, \quad (9)$$

which monotonically increases with $|\alpha|$.

As mentioned previously, the case $k = 2$ ($\varphi_k = \pi$) corresponds to the creation of the vacuum with squeezed

quadratures. We find high (here and henceforth meaning above 99%) overlap with

$$|\Psi_2\rangle = S(z)|0\rangle = e^{-\frac{1}{2}(z(\hat{a}^\dagger)^2 - z^*(\hat{a})^2)}|0\rangle \quad (10)$$

by numerically searching over N and z , where $S(z)$ is the ordinary squeezing operator with complex squeezing parameter z . In general, for $k \geq 2$, the states created by this protocol are very close to the analytic states

$$|\Psi_k\rangle = S^{(k)}(z)|0\rangle = e^{-\frac{1}{2}(z(\hat{a}^\dagger)^k - z^*(\hat{a})^k)}|0\rangle, \quad (11)$$

where $S^{(k)}(z)$ is called a generalized squeezing operator with complex parameter z [20, 21]. The first three operators of this class are

$$S^{(0)}(z) = e^{-\text{Im}[z]\mathbb{I}} \quad (12)$$

$$S^{(1)}(z) = D\left(-\frac{z}{2}\right) \quad (13)$$

$$S^{(2)}(z) = S(z) \quad (14)$$

where $S(z)$ is the squeezing operator of equation (10).

Consider the final state after displacing to the origin ρ_k . To find z , we impose the condition that ρ_k and $|\Psi_k\rangle$ have the same average photon number by setting

$$\langle \Psi_k | \hat{a}^\dagger \hat{a} | \Psi_k \rangle = \langle \hat{a}^\dagger \hat{a} \rangle_{\rho_k}. \quad (15)$$

This results in excellent fidelity between the states, quantified by

$$F[\rho_k] \equiv \text{Tr}[\rho_k |\Psi_k\rangle \langle \Psi_k|]. \quad (16)$$

In fact, for $|\alpha| = 3$ and $k = 2, 3, 4$ the fidelity is $> 99\%$ in each case. The Husimi Q functions of these states are shown in figures 2 (a), (c) and (e).

To verify that these states are indeed nonclassical, a suitable nonclassicality witness can be used. For this purpose, we use the entanglement potential of [22], defined by

$$\text{EP}[\rho] \equiv \log_2 \|\sigma_\rho^{TA}\|_1, \quad (17)$$

where $\sigma_\rho = U_{\text{BS}}(\rho \otimes |0\rangle\langle 0|)U_{\text{BS}}^\dagger$, for U_{BS} the unitary transformation of a 50:50 beam splitter, and σ_ρ^{TA} is the partial transpose of σ_ρ [23]. A nonzero entanglement potential indicates that the state ρ is nonclassical, and indeed, as can be calculated numerically, the states created by the generalized protocol all have nonzero entanglement potential (see supplementary material).

Since the displacement stage of each step ensures the average photon number of each state ρ_j is on the order of $|\alpha|^2$, each P_N^j can be quite large, and as a result, the generalized protocol will have a significant probability of success. For example, for the modest $|\alpha| = 3$ of the states shown in figure 2, the success probabilities are all $\approx 85\%$. Furthermore, the success probability will scale monotonically with $|\alpha|$, and so can be increased by increasing the initial input state power.




FIG. 2: This figure shows the Husimi Q functions for the states created by the generalized protocol, figures (a), (c) and (e), and those created by one additional photon detection after displacement to the origin, figures (b), (d) and (f).

This protocol can be generalized even further to create additional nonclassical states, including the squeezed odd Schrödinger cat state, with only a single additional detection required. This extension to the generalized protocol is similar to that proposed in [6] to prepare Schrödinger cat-like states, but does not require a beam splitter or photon number resolving detectors.

If we remove one more photon from the final state ρ_k , the resulting state has the form

$$\rho'_k = \frac{\hat{B}\rho_k\hat{B}^\dagger}{P_1^k}, \quad (18)$$

where P_1^k is the probability of a single photon being detected from state ρ_k . This procedure produces states of high overlap with

$$|\Psi'_k\rangle = S^{(k)}(z) \sum_{j=0}^{k-1} |\beta e^{ij\varphi_k}\rangle, \quad (19)$$

where now both z and β must be found numerically. These states all have non-zero entanglement potential, and are therefore nonclassical states of microwave radiation (see supplementary material).

For $k = 2$, this additional detection will create a state very close to a squeezed odd Schrödinger cat state

$$|\Psi'_2\rangle = S^{(2)}(z) (|\beta\rangle - |-\beta\rangle). \quad (20)$$

For $k = 3$ and $k = 4$, we have

$$|\Psi'_3\rangle = S^{(3)}(z) \left(|\beta\rangle + \left| \beta e^{\frac{2\pi}{3}} \right\rangle + \left| \beta e^{\frac{4\pi}{3}} \right\rangle \right) \quad (21)$$

$$|\Psi'_4\rangle = S^{(4)}(z) (|\beta\rangle + |i\beta\rangle + |-\beta\rangle + |-i\beta\rangle). \quad (22)$$

The $k = 3$ state is the state formed when the operator $S^{(3)}(z)$ is applied to a Voodoo cat state [9]. The $k = 4$ state is the operator $S^{(4)}(z)$ applied to a coherent superposition of four out of phase coherent states, known as a compass state, which is known to have favourable decoherence properties [13, 24].

The Husimi Q functions of these squeezed cat-like states (for $|\alpha| = 3$) are shown in figure 2, all of which have greater than 99% fidelity with (20), (21), and (22). For the $k = 4$ case, it was necessary to slightly modify the shifts α_1 and α_3 used in the generalized protocol by a small increment $-\delta|e^{i(j\varphi_k + \varphi_\alpha)}$ to achieve high fidelity. For the $|\alpha| = 3$ case shown, $|\delta| \approx 0.1$.

The probability of successful generation of these squeezed nonclassical states is

$$\text{Prob}(\text{success})' = P_1^k \prod_{j=0}^{k-1} P_N^j. \quad (23)$$

Unfortunately, these P_1^k are often very small, and for the squeezed cat-like states shown in figures 2 (b), (d) and (f) this results in a success probability less than 10%. However, as P_1^k grows monotonically with $|\alpha|$, the success probability improves with higher power coherent pulses.

In regards to experimental implementation of this protocol with JPMs, superconducting microwave resonators are currently fabricated with Q-factors approaching 10^7 , which in the microwave regime will lead to cavity lifetimes on the order of 10^5 ns [25]. Thus, when JPMs with short T_2 are used [16], we can conservatively expect that as many as 10^2 to 10^3 measurements can be performed in the lifetime of the cavity. Practically, space requirements on a chip require a fast reset strategy for the JPMs, which is currently being developed [26].

In conclusion, we have shown how a combination of strongly coupled photon counting and coherent displacement can be used to create nonclassical states of radiation with high probability. This protocol can be realized in circuit QED using Josephson photomultipliers.

We acknowledge instructive discussions with F.W. Strauch and R. McDermott. Research supported by DARPA within the QuEST program and NSERC Discovery grants. LCGG was supported by the Ontario Graduate Scholarship Program.

SUPPLEMENTARY MATERIAL




FIG. 3: This figure shows entanglement potential (EP) of the states created by the generalized protocol (for $k = 2, 3, 4$) as a function of the number of detections, N , performed at each step in the protocol. The generalized squeezed states are the solid curves, and the dotted curves correspond to the cat-like states. These states have $|\alpha| = 3$.

The entanglement potential is not only a nonclassicality witness but is in fact also a nonclassicality measure. Here we examine how the nonclassicality of the states changes with the number of detections at each step of the generalized protocol. Figure 3 shows the entanglement potential of the generalized squeezed states and the cat-like states (for $k = 2, 3, 4$) as a function of the number of detections at each step. As can be seen, for the squeezed state ($k = 2$), the maximum value of EP occurs around the optimal value of N determined in the main text. In addition, for each state shown in figure 3, the value of N which gave the highest fidelity with the ideal state corresponds to a near maximum value of the entanglement potential.

[1] E. Knill, R. Laflamme, and G. Milburn, *Nature* **409**, 46 (2001).
 [2] S. Braunstein and P. van Loock, *Rev. Mod. Phys.* **77**, 513 (2005).
 [3] N. Gisin, R. Ribordy, W. Tittel, and H. Zbinden, *Rev. Mod. Phys.* **74**, 145 (2002).

[4] A. I. Lvovsky and M. G. Raymer, *Rev. Mod. Phys.* **81**, 299 (2009).
 [5] A. Ourjoumtsev, R. Tualle-Broui, J. Laurat, and P. Grangier, *Science* **312**, 83 (2006).
 [6] N. Dakna, T. Anhut, T. Opatrny, L. Knöll, and D.-G. Welsch, *Phys. Rev. A* **55**, 3184 (1997).

- [7] M. K. Akhlaghi, A. H. Majedi, and J. S. Lundeen, *Optics Express* **19**, 21305 (2011).
- [8] M. Hofheinz, E. M. Weig, M. Ansmann, R. C. Bialczak, E. Lucero, M. Neeley, A. D. O'Connell, H. Wang, J. M. Martinis, and A. N. Cleland, *Nature* **454**, 310 (2008).
- [9] M. Hofheinz, H. Wang, M. Ansmann, R. C. Bialczak, E. Lucero, M. Neeley, A. D. O'Connell, D. Sank, J. Wenner, J. M. Martinis, et al., *Nature* **459**, 546 (2009).
- [10] M. A. Castellanos-Beltran, K. D. Irwin, G. C. Hilton, L. R. Vale, and K. W. Lehnert, *Nat Phys* **4**, 929 (2008).
- [11] C. Eichler, D. Bozyigit, C. Lang, M. Baur, L. Steffen, J. M. Fink, S. Filipp, and A. Wallraff, *Phys. Rev. Lett.* **107**, 113601 (2011).
- [12] C. Wilson, G. Johansson, A. Pourkabirian, J. Johansson, T. Duty, F. Nori, and P. Delsing, *Nature* **479**, 376 (2011).
- [13] Z. Leghtas, G. Kirchmair, B. Vlastakis, M. Devoret, R. Schoelkopf, and M. Mirrahimi (2012), arXiv:1205.2401.
- [14] J. M. Raimond, C. Sayrin, S. Gleyzes, I. Dotsenko, M. Brune, S. Haroche, P. Facchi, and S. Pascasio, *Phys. Rev. Lett.* **105**, 213601 (2010).
- [15] Y.-F. Chen, D. Hover, S. Sendelbach, L. Maurer, S. T. Merkel, E. J. Pritchett, F. K. Wilhelm, and R. McDermott, *Phys. Rev. Lett.* **107**, 217401 (2011).
- [16] L. C. G. Govia, E. J. Pritchett, S. T. Merkel, D. Pineau, and F. K. Wilhelm, *Phys. Rev. A* **86**, 032311 (2012).
- [17] A. Poudel, R. McDermott, and M. Vavilov, arXiv:1201.2990.
- [18] B. Peropadre, G. Romero, G. Johansson, C. Wilson, E. Solano, and J. Garcia-Ripoll, *Phys. Rev. A* **84**, 063834 (2011).
- [19] G. B. Arfken, H. J. Weber, and F. E. Harris, *Mathematical Methods for Physicists: A Comprehensive Guide* (Academic Press, 2012), 7th ed.
- [20] S. L. Braunstein and R. I. McLachlan, *Phys. Rev. A* **35**, 1659 (1987).
- [21] S. L. Braunstein and C. M. Caves, *Phys. Rev. A* **42**, 4115 (1990).
- [22] J. K. Asbóth, J. Calsamiglia, and H. Ritsch, *Phys. Rev. Lett.* **94**, 173602 (2005).
- [23] G. Vidal and R. F. Werner, *Phys. Rev. A* **65**, 032314 (2002).
- [24] W. H. Zurek, *Nature* **412**, 712 (2001).
- [25] A. Megrant, C. Neill, R. Barends, B. Chiaro, Y. Chen, L. Feigl, J. Kelly, E. Lucero, M. Mariantoni, P. J. J. O'Malley, et al., *Applied Physics Letters* **100**, 113510 (pages 4) (2012).
- [26] R. McDermott (2012), private communication.
- [27] for which the analytic expression is cumbersome and not instructive but has the same limiting values as P_N for $|\alpha| \rightarrow \infty$